

One-parameter Darboux transformations in Thermodynamics

Haret C. Rosu¹

Applied Mathematics Dept., IPICyT, Apdo Postal 3-74 Tangamanga,
San Luis Potosí, SLP, Mexico

Abstract

The quantum oscillator thermodynamic actions are the conjugate intensive parameters for the frequency in any frequency changing process. These oscillator actions fulfill simple Riccati equations. Interesting Darboux transformations of the fundamental Planck and pure vacuum actions are discussed here in some detail. It is shown that the one-parameter “Darboux-Transformed-Thermodynamics” refers to superpositions of boson and fermion excitations of positive and negative absolute temperature, respectively. A Darboux generalization of the fluctuation-dissipation theorem is also briefly sketched.

Using the Planck energy distribution of a quantum oscillator one can define an action function

$$f_P(x) = \frac{U_P}{\omega} = \frac{\hbar}{2} + \frac{\hbar}{\exp(\hbar x) - 1} = \frac{\hbar}{2} \coth\left(\frac{\hbar x}{2}\right), \quad (1)$$

where $x = \beta\omega$. This action plays the role of a generalized force in any process of frequency change performed by the oscillator.

The point now is that the function $f_P(x)$ fulfills as particular solution the following Riccati equation

$$\frac{df}{dx} + f^2 = \left(\frac{\hbar}{2}\right)^2. \quad (2)$$

As a matter of fact, the constant vacuum action $f_V = \frac{\hbar}{2}$ is also a solution of the same Riccati equation, whereas the pure thermal action $f_T = \frac{\hbar}{\exp(\hbar x) - 1}$ is a particular solution of the following equation

$$\frac{df}{dx} + \hbar f + f^2 = 0, \quad (3)$$

¹E-mail: hcr@ipicyt.edu.mx

Contribution at the AIP 1st International Conference on Quantum Limits to the Second Law, San Diego, CA, (main organizer: D.P. Sheehan), end of July 2002, tentative publication schedule: Dec. 2002.

which is again a particular type of Riccati equation (a Bernoulli equation). Since the Riccati equation is a main ingredient in supersymmetric quantum mechanics [1] we would like to present here some thermodynamic consequences of using methods belonging to the area which focus on one-parameter Darboux transformations of thermodynamic actions.

In supersymmetric quantum mechanics [2], which is a simple application of Darboux transformations [1], one usually starts with a known particular solution of a Riccati equation without linear term

$$\frac{df_p}{dx} + f_p^2 = V_1(x) \quad (4)$$

that we call the bosonic Riccati equation. We are not interested here in a so-called factorization constant that can be placed at the right hand side. The function V_1 is an exactly solvable potential for the (“imaginary-time”) Schrödinger equation at zero energy

$$\left(\frac{d}{dx} + f_p\right) \left(\frac{d}{dx} - f_p\right) w_b = \frac{d^2 w_b}{dx^2} - V_1(x) w_b = 0 . \quad (5)$$

The particular solution w_b is usually called a bosonic zero mode. It is connected to the Riccati solution through $f_p = \frac{1}{w_b} dw_b/dx$. Next, changing the sign of the first derivative in Eq. (4) one calculates the outcome $V_2(x)$ using the same Riccati solution

$$-\frac{df_p}{dx} + f_p^2 = V_2(x) . \quad (6)$$

We call the latter equation the fermionic Riccati equation. The function $V_2(x)$ is known as the supersymmetric partner of the initial potential $V_1(x)$. The corresponding zero mode fulfills

$$\left(\frac{d}{dx} - f_p\right) \left(\frac{d}{dx} + f_p\right) w_f = \frac{d^2 w_f}{dx^2} - V_2(x) w_f = 0 . \quad (7)$$

On the other hand, already in 1984, Mielnik [3] studied the ambiguity of the factorization of the Schrödinger equation for the oscillator that led him to the general (one-parameter-dependent) solution of the companion Riccati equation for that case. In other words, one looks for the general solution [1]

$$-\frac{df_g}{dx} + f_g^2 = -\frac{df_p}{dx} + f_p^2 = V_2(x) . \quad (8)$$

The latter equation can be solved for f_g by employing the Bernoulli ansatz $f_g(x) = f_p(x) - \frac{1}{v(x)}$, where $v(x)$ is an unknown function [4]. One obtains for the function $v(x)$ the following Bernoulli equation

$$\frac{dv(x)}{dx} + 2v(x) f_p(x) = 1 \quad (9)$$

that has the solution

$$v(x) = \frac{\mathcal{I}_{0b}(x) + \lambda}{w_b^2(x)}, \quad (10)$$

where $\mathcal{I}_{0b}(x) = \int_0^x w_b^2(y) dy$, and we consider λ as a positive integration constant that is employed as a free parameter.

Thus, the general fermionic Riccati solution is a one-parameter function $f_g(x; \lambda)$ of the following form

$$f_g(x; \lambda) = f_p(x) - \frac{d}{dx} [\ln(\mathcal{I}_{0b}(x) + \lambda)] = \frac{d}{dx} \left[\ln \left(\frac{w_b(x)}{[\mathcal{I}_{0b}(x) + \lambda]} \right) \right] = \frac{d}{dx} \ln w_b(x; \lambda) \quad (11)$$

where $w_b(x; \lambda) = \frac{w_b(x)}{\mathcal{I}_{0b}(x) + \lambda}$. The range of the λ parameter is conditioned by $\mathcal{I}_0(x) + \lambda \neq 0$ in order to avoid singularities. This is a well-known restriction [3]. According to the supersymmetric construction one can use this general fermionic Riccati solution to calculate a one-parameter family of bosonic potentials as follows

$$\frac{df_g}{dx} + f_g^2 = V_{1,g}, \quad (12)$$

where

$$V_{1,g} = V_1 - 2 \frac{d^2}{dx^2} \ln(\mathcal{I}_{0b}(x) + \lambda) \quad (13)$$

enters the linear equation

$$\left(\frac{d}{dx} + f_g \right) \left(\frac{d}{dx} - f_g \right) w_b(x; \lambda) = \frac{d^2 w_b(x; \lambda)}{dx^2} - V_{1,g} w_b(x; \lambda) = 0. \quad (14)$$

In the limit $\lambda \rightarrow \infty$, Eq. (12) goes into Eq. (4) because $f_g \rightarrow f_p$ and $V_{1,g} \rightarrow V_1$.

One can think of Eq. (12) as a generalization of the thermodynamic Riccati equation (2). Of interest are the one-parameter oscillator actions f_g

rather than the ‘potentials’ $V_{1,g}$. Since in supersymmetric quantum mechanics $V_{1,g}$ are the general Darboux-transformed potentials, we shall call the functions f_g as Darboux-transformed actions.

Going now back to Eq. (2) and using $f = w'/w$, where the prime denotes the derivative with respect to x , one gets the linear equation

$$w'' - \left(\frac{\hbar}{2}\right)^2 w = 0 , \quad (15)$$

having the particular zero-mode solution $w_a = W_a \sinh(\frac{\hbar}{2}x)$. Thus, the general Riccati solution is a one-parameter function $f_{gP}(x; \lambda)$ of the following form

$$f_{gP}(x; \lambda) = f_P(x) - \frac{d}{dx} \left[\ln(\mathcal{I}_{0a}(x) + \lambda) \right] = \frac{d}{dx} \left[\ln \left(\frac{w_a(x)}{[\mathcal{I}_{0a}(x) + \lambda]} \right) \right]. \quad (16)$$

For the vacuum case the particular Riccati solution is the vacuum action $f_p = f_V = \frac{\hbar}{2}$. The corresponding zero mode is $w_V \propto e^{\hbar x/2}$. The usual entropy of the vacuum fluctuations is zero whereas the modified entropy based on f_{gV} has a kink-like behavior between the $\frac{\hbar}{2}$ (bosonic) solution and the $-\frac{\hbar}{2}$ (fermionic) solution.

One can also use the general zero-mode $w_g = Ae^{\hbar x/2} + Be^{-\hbar x/2}$ as solution in Eq. (5). The Planck action corresponds to $A = -B = \frac{1}{2}$ (antisymmetric zero-mode), while the vacuum case to $A = \text{arbitrary}$ and $B = 0$. A very interesting case corresponds to the symmetric zero-mode, $A = B = \frac{1}{2}$, $w_s = W_s \cosh(\frac{\hbar}{2}x)$, since if we trace back to the action we get

$$f_s = -\frac{\hbar}{2} + \frac{\hbar}{\exp(-\hbar x) + 1} , \quad (17)$$

i.e., a Fermi-Dirac action for negative x . We have shown in a previous paper that the λ parameter is equivalent to the quotient A/B [5]. Thus, the general Riccati solution introduces effects of the second linear independent solution. The problem then turns into a subtle interpretation of the mathematical results. We have found at least one possible physical significance. For the Planck case, the second linear independent zero-mode is exactly the cosh function and the corresponding action is the aforementioned Fermi-Dirac action of negative x . We attach the minus sign to the temperature parameter

in $x = \omega/T$ rather than to ω and recall that the issue of negative absolute (spin) temperatures first appeared in Physics in 1951 when Purcell and Pound were able to produce sudden reversals of the direction of an external magnetic field applied to a crystal of LiF [6]. Since then many other experiments with negative temperatures have been devised in nuclear spin systems and the corresponding ‘violations’ of the second law of thermodynamics were a subject of discussion [7]. Thus, the one-parameter Darboux transformations of the Planck action are a way of introducing the effect of a Fermi-Dirac action of negative absolute temperature.

Another interesting application refers to dissipative *RLC* systems where the oscillator action enters Nyquist-Johnson spectral power noise distributions of the type (fluctuation-dissipation theorem [8])

$$P(\omega, \beta) = \frac{\omega}{\pi} R(\omega, \beta) f_P . \quad (18)$$

One can think of the corresponding generalization

$$P(\omega, \beta; \lambda) = \frac{\omega}{\pi} R(\omega, \beta) f_g(\beta\omega; \lambda) \quad (19)$$

and hope to study even in simple experiments the significance of this Darboux generalization of the fluctuation-dissipation theorem.

In conclusion, Planck’s thermodynamic oscillator action can be generalized to a one parameter Darboux family of actions. We also consider the bosonic vacuum case separately in the same way. The Planck action and the pure vacuum case correspond to the asymptotic limit of the Darboux parameter $\lambda \rightarrow \infty$. In the Planck case, all the other λ cases correspond to a system made of bosons at temperature T interacting with an equal system of fermions at temperature $-T$. In the vacuum case, the $\lambda \neq \infty$ cases describe the interaction of the bosonic and fermionic vacua. Using the procedure proposed by Arnaud, Chusseau and Philippe [9] we have calculated the efficiencies of the ideal oscillator Carnot cycles based on the Darboux-modified Planck and vacuum entropies in a previous work [10]. Systems of negative Kelvin temperatures are hotter than those of positive T [7] and therefore they always represent the hot bath. In real, dissipative cases, the same type of generalization is suggested for the fluctuation-dissipation theorem. Finally, we mention that a multiple-parameter Darboux generalization is also possible [11]. The experimental nature of the λ parameter is an open issue.

References

- [1] H.C. Rosu, *Short survey of Darboux transformations*, in Proc. “Symmetries in Quantum Mechanics and Quantum Optics”, Burgos, Spain, Sept. 21-24, 1998, Eds. A. Ballesteros et al., (Serv. de Publ., Univ. Burgos, Burgos, Spain, 1999) pp. 301-315, quant-ph/9809056.
- [2] E. Witten, Nucl. Phys. B **188**, 513 (1981).
- [3] B. Mielnik, J. Math. Phys. **25**, 3387 (1984); For a recent application to cosmology, see H.C. Rosu, Mod. Phys. Lett. A **15**, 979 (2000), gr-qc/0003108.
- [4] D. Zwillinger, *Handbook of Differential Equations* (Academic Press, New York, 1989).
- [5] L.J. Boya, H. Rosu, A.J. Seguí-Santonja, F.J. Vila, Nuovo Cimento B **113**, 409 (1998), quant-ph/9711059.
- [6] E.M. Purcell and R.V. Pound, Phys. Rev. **81**, 279 (1951).
- [7] N.F. Ramsey, Phys. Rev. **103**, 20 (1956). See also, N. Bloembergen, Am. J. Phys. **41**, 325 (1973).
- [8] H. Nyquist, Phys. Rev. **32**, 110 (1928).
- [9] J. Arnaud, L. Chusseau, and F. Philippe, Eur. J. Phys. **23**, 489 (2002), physics/0105048.
- [10] H.C. Rosu, F. Aceves de la Cruz, Physica Scripta **65**, 377 (2002), quant-ph/0107043
- [11] H.C. Rosu, Int. J. Theor. Phys. **39**, 105 (2000), quant-ph/9904007.